

No evidence for widespread bird declines in protected South American forests

**Philip C Stouffer · Kristina L. Cockle · Alexandre Aleixo · Juan I. Areta ·
Juan Mazar Barnett · Alejandro Bodrati · Carlos Daniel Cadena ·
Adrián S. Di Giacomo · Sebastian K. Herzog · Peter Hosner · Erik I. Johnson ·
Luciano N. Naka · César Sánchez**

Received: 19 January 2010 / Accepted: 20 June 2011 / Published online: 23 July 2011
© Springer Science+Business Media B.V. 2011

As ornithologists working with birds in South America, we were intrigued by the title of a recent paper in *Climatic Change*, ‘Are bird populations in tropical and subtropical forests of South America affected by climate change?’ (Nores 2009). The author claims that vast areas of South American forests have become ‘practically ornithological deserts,’ presumably due to some unspecified effect of climate change. However, the bird data reported grossly underestimate current bird diversity in these forests. Moreover, the methods employed are unacceptable in modern ornithology, and the conclusions of the

P. C. Stouffer (✉) · E. I. Johnson
School of Renewable Natural Resources, Louisiana State University Agricultural Center,
Baton Rouge, LA 70803, USA
e-mail: pstouffer@lsu.edu

P. C. Stouffer · E. I. Johnson
Projeto Dinâmica Biológica de Fragmentos Florestais, Instituto Nacional de Pesquisas da Amazônia
and Smithsonian Tropical Research Institute, CP 478, Manaus, AM 69060, Brazil

K. L. Cockle
Center for Applied Conservation Research, Department of Forest Sciences,
University of British Columbia, Vancouver, BC V6T 1Z4, Canada

A. Aleixo
Coordenação de Zoologia, MCT/Museu Paraense Emílio Goeldi, Belém, Pará, Brazil

J. I. Areta
CICyTTP-CONICET, Materi & España, Diamante (3105) Entre Ríos, Argentina

J. M. Barnett
Plaza 2407 2*A, (1430) Buenos Aires, Argentina

A. Bodrati
Proyecto Selva de Pino Paraná, San Pedro, Misiones, Argentina

C. D. Cadena
Departamento de Ciencias Biológicas, Universidad de los Andes, Bogotá, Colombia

paper are deeply flawed. The paper misrepresents the state of bird diversity in South American forests, and provides no insights on possible effects of climate change on birds in these forests.

The most glaring defect of the paper is the quantitative data on richness and abundance of birds. Apparently, Nores reported only the birds he identified visually, although acoustic surveys have long been recognized as absolutely essential to quantify bird diversity in tropical forest (e.g., Terborgh et al. 1990). While Nores (2009) reports “no more than 15 species and 18 individuals in one day” for some Amazon forest sites, our single-day lists based on visual and auditory samples typically include some 70–100 species in large forest tracts. At Parque Nacional Iguazú, a site emphasized by Nores, Hosner found about twice as many species/day in 2005. In visits to Iguazú from 2007 to 2009, Areta, Bodrati and Cockle detected 77 to 174 species/day, including daily records of every species that Nores (2009) cited as absent (Table 3 in Nores 2009). Recent research also shows high species richness and abundance from other South American tropical and subtropical forests, including some sites mentioned in Nores (2009). For example, Herzog (2008) reported 131–190 species from three Bolivian Yungas locations. Johnson et al. (2011) found 228 species in 2008 in a 100-ha forest plot north of Manaus, Brazil, with over 100 species recorded/day. Bodrati et al. (2010) found 312 species in 600-ha Parque Provincial Cruce Caballero, 180 km southeast of Iguazú. Clearly, avian diversity in tropical forests is grossly underestimated by single-day lists based only on birds identified visually.

Even if the single-day lists reported by Nores (2009) had included a reasonable proportion of the detectable birds, raw lists provide an unsuitable metric for analyzing spatial or temporal change in species richness or abundance (Remsen 1994). A major focus of ecological research has been to develop robust estimates for these parameters from field data inevitably based on incomplete sampling and imperfect detection (e.g., Seber 1986; Nichols et al. 1998; Buckland et al. 2001; Colwell 2009). Not surprisingly, replication or subsampling become the key for drawing inferences, particularly separating lack of detection from actual absence (Thompson 2002; Herzog et al. 2002; MacKenzie and Royle 2005). Nores completely ignored these sampling and analytical issues, even though they are increasingly applied to Neotropical bird communities (e.g. Blake 2007; Herzog and Kessler 2006).

Finally, even if the field lists had been plausible and the samples had been subjected to a procedure to produce meaningful estimates of abundance or species richness, the paper presents no analysis to support the claim that there has actually been a temporal change. Given that a major focus of the climate change literature is temporal changes of physical or biological variables, the absence of any temporal analysis (or even estimate of uncertainty) seriously reduces the value of this work. To be fair, the paper only states that the purported

A. S. Di Giacomo

Departamento de Ecología, Genética y Evolución, FCEN-Universidad de Buenos Aires,
Pabellón II Ciudad Universitaria C1428EHA, Buenos Aires, Argentina

S. K. Herzog

Asociación Armonía - BirdLife International, Santa Cruz de la Sierra, Bolivia

P. Hosner

Natural History Museum & Biodiversity Research Center,
University of Kansas, Lawrence, KS 66044, USA

L. N. Naka · C. Sánchez

Museum of Natural Science and Department of Biological Sciences, Louisiana State University,
Baton Rouge, LA 70803, USA

change in bird diversity ‘could be associated with climate change,’ as also evidenced by the question mark in the title. Clearly, however, the emphasis of the paper is that climate change, through some unspecified mechanism, has led to widespread crashes in bird diversity in the most species-rich forests in the world. We see no evidence to support that claim, neither from the paper nor from our own recent fieldwork.

We do not deny that climate change could be affecting South American tropical and subtropical birds. Indeed, we encourage research to elucidate its mechanisms and effects. We agree with Nores (2009) that continued monitoring at sites with a history of ornithological research could be especially useful. Based on results from other continents, the most obvious places to look for effects of climate change would be phenological effects in breeding and migration (e.g. Parmesan 2006; Gordo 2007). Elevational shifts or other changes in distribution might also be revealed by careful study (e.g. Pounds et al. 1999). As a threat to Neotropical birds, however, we believe that climate change presents a far less immediate problem than forest loss. South America loses about 2.5 million ha of humid tropical forest per year to deforestation (Achard et al. 2002), with strong and direct impacts on forest birds at the local and regional scale (e.g. Ribon et al. 2003; Lees and Peres 2006; Stouffer et al. 2006, 2009; Giraudo et al. 2008). The most important step to conserve forest bird communities in South America remains the protection of large tracts of native forest. To be relevant for conservation of South American forest birds, research on climate change needs to take into account the concurrent and possibly synergistic effects of forest loss, fragmentation, and bird species’ distributions (Travis 2003; Jetz et al. 2007; Laurance and Useche 2009; Ribeiro et al. 2009). We encourage quality field research, modern analytical methods, and rigorous peer-review to identify effects of climate change on tropical birds. We see nothing in Nores (2009) that advances our understanding in this important research area.

Acknowledgements Funding for the research in Manaus was provided by the US National Science Foundation (LTREB-0545491). This is publication 574 of the BDFFP INPA/STRI Technical Series and 21 of the Amazonian Ornithology Technical Series of the INPA Collections Program. The manuscript was approved by the Director of the Louisiana State University Agricultural Center as manuscript number 2011-241-6154.

References

- Achard F, Eva HD, Stibig HJ, Mayaux P, Gallego J, Richards T, Malingreau JP (2002) Determination of deforestation rates of the world’s humid tropical forests. *Science* 297:999–1002
- Blake JG (2007) Neotropical forest bird communities: a comparison of species richness and composition at local and regional scales. *Condor* 109:237–255
- Bodrati A, Cockle K, Segovia JM, Roesler I, Areta JI, Jordan E (2010) La avifauna del Parque Provincial Cruce Caballero, Provincia de Misiones, Argentina. *Cotinga* 32:41–64
- Buckland ST, Anderson DR, Burnham KP, Laake JL, Borchers DL, Thomas L (2001) Introduction to distance sampling. Estimating abundance of biological populations. Oxford University Press, New York
- Colwell RK (2009) EstimateS 8.2 user’s guide. <http://viceroy.eeb.uconn.edu/estimates>, accessed 6 January 2010
- Giraudo AR, Matteucci SD, Alonso J, Herrera J, Abramson RR (2008) Comparing bird assemblages in large and small fragments of the Atlantic Forest hotspots. *Biodivers Conserv* 17:1251–1265
- Gordo O (2007) Why are bird migration dates shifting? A review of weather and climate effects on avian migratory phenology. *Clim Res* 35:37–58
- Herzog SK (2008) Primer relevamiento ornitológico en la Cordillera Mosetenes, con una comparación latitudinal de comunidades de aves en los Yungas bajos de Bolivia/First ornithological survey of Cordillera Mosetenes, with a latitudinal comparison among lower Yungas bird communities in Bolivia. *Rev Bol Biol, Ecol y Cons Amb* 23:59–71
- Herzog SK, Kessler M (2006) Local vs. regional control on species richness: a new approach to test for competitive exclusion at the community level. *Glob Ecol Biogeogr* 15:163–172

- Herzog SK, Kessler M, Cahill TM (2002) Estimating species richness of tropical bird communities from rapid assessment data. *Auk* 119:749–769
- Jetz W, Wilcove DS, Dobson AP (2007) Projected impacts of climate and land-use change on the global diversity of birds. *PLoS Biol* 5:1211–1219
- Johnson EI, Stouffer PC, Vargas CF (2011) Diversity, biomass, and trophic structure of a central Amazonian rainforest bird community. *Rev Bras Ornith* 19:1–16
- Laurance WF, Useche DC (2009) Environmental synergisms and extinctions of tropical species. *Conserv Biol* 23:1427–1437
- Lees AC, Peres CA (2006) Rapid avifaunal collapse along the Amazonian deforestation frontier. *Biol Conserv* 133:198–211
- Mackenzie DI, Royle JA (2005) Designing occupancy studies: general advice and allocating survey effort. *J Appl Ecol* 42:1105–1114
- Nichols JD, Boulinier T, Hines JE, Pollock KH, Sauer JR (1998) Inference models for spatial variation in species richness and community composition when not all species are detected. *Conserv Biol* 12:1399–1404
- Nores M (2009) Are bird populations in tropical and subtropical forests of South America affected by climate change? *Clim Chang* 97:543–551
- Parnesan C (2006) Ecological and evolutionary responses to recent climate change. *Annu Rev Ecol Evol Syst* 37:637–669
- Pounds JA, Fogden MPL, Campbell JH (1999) Biological response to climate change on a tropical mountain. *Nature* 398:611–615
- Remsen JV Jr (1994) Use and misuse of bird lists in community ecology and conservation. *Auk* 111:225–227
- Ribeiro MC, Metzger JP, Martensen AC, Ponzoni FJ, Hirota MM (2009) The Brazilian Atlantic forest: how much is left, and how is the remaining forest distributed? Implications for conservation. *Biol Conserv* 142:1141–1153
- Ribon R, Simon JE, De Mattos GT (2003) Bird extinctions in Atlantic forest fragments of the Vicosa region, southeastern Brazil. *Conserv Biol* 17:1827–1839
- Seber GAF (1986) A review of estimating animal abundance. *Biometrics* 42:267–292
- Stouffer PC, Bierregaard RO Jr, Strong C, Lovejoy TE (2006) Long-term landscape change and bird abundance in Amazonian rainforest fragments. *Conserv Biol* 20:1212–1223
- Stouffer PC, Strong C, Naka LN (2009) Twenty years of understorey bird extinctions from Amazonian rain forest fragments: consistent trends and landscape-mediated dynamics. *Divers Distrib* 15:88–97
- Terborgh J, Robinson SK, Parker TA III, Munn CA, Pierpont N (1990) Structure and organization of an Amazonian forest bird community. *Ecol Monogr* 60:213–238
- Thompson WL (2002) Towards reliable bird surveys: accounting for individuals present but not detected. *Auk* 119:18–25
- Travis JMJ (2003) Climate change and habitat destruction: a deadly anthropogenic cocktail. *Proc R Soc Lond B* 270:467–473